

PRIVILEGED AND CONFIDENTIAL
ATTORNEY WORK PRODUCT

COVINGTON & BURLING DRAFT

May 4, 1994

TOBACCO INGREDIENT PYROLYSIS AND TRANSFER CONTRIBUTIONS
TO CIGARETTE MAINSTREAM SMOKE

ABSTRACT

This report summarizes information about tobacco leaf composition and the contribution of tobacco leaf components to the composition of mainstream cigarette smoke, compares cigarette additive composition with tobacco leaf composition, and describes the relationships between cigarette additives and cigarette mainstream smoke composition.

A significant amount of research on the composition of tobacco leaf, tobacco smoke and the contribution of leaf components to cigarette smoke has been conducted over the past 40 years. The research has resulted in the development of a vast amount of information on the contribution of leaf components to cigarette smoke. The information on leaf components can be used to assess the contribution of tobacco additives to mainstream smoke.

Many of the major additives used in cigarettes replace naturally occurring components of tobacco which are lost in the curing process. Additives which are identical to leaf components are sugars, starches, organic acids and their

2029063045

salts, lipids, proteins and amino acids, and many of the volatile flavor ingredients. Other major additives which are not tobacco identical are essentially similar in composition to natural leaf components. These include materials such as cocoa, licorice, honey, corn syrup, sugar alcohols, and fruit juice concentrates. By virtue of their compositional and structural similarity to tobacco leaf components, the additives typically used in cigarettes are expected to exhibit similar transfer behavior or to contribute substantially the same types of pyrolysis products to mainstream smoke. Other substances that are used as processing aids, such as ethyl alcohol and carbon dioxide, do not contribute significantly to mainstream smoke composition because virtually none of these materials is present in the tobacco product after manufacture.

BACKGROUND

Cigarette manufacturers and other organizations have conducted and sponsored a significant amount of research into the composition of tobacco leaf and tobacco smoke. This research provides comprehensive information on the contribution of tobacco leaf components to mainstream cigarette smoke. Natural leaf components representing the majority of tobacco leaf dry weight are carbohydrates, proteins, amino acids, organic acids, phenolics, nicotine, waxes and oils. Many of the major tobacco additives belong to

2029063046

these same classes of compounds. The structural similarities between natural leaf constituents and the substances added to tobacco allow conclusions to be drawn on the relationship between cigarette blend components and their contribution to mainstream cigarette smoke.

The vast majority (by number, not poundage) of ingredients added to tobacco -- the flavor ingredients -- are volatile compounds by virtue of their low boiling points or ease of sublimation. Examples of additives that fall in this category are menthol, ethyl acetate, and vanillin. This category also includes certain volatile constituents of essential oils such as nutmeg/mace, anise/anise oil and star anise/star anise oil. Because of their physical and chemical properties, these volatile ingredients do not undergo pyrolysis to any significant extent. Rather, they transfer unchanged from tobacco to the mainstream smoke at an average rate of 12% to 15% of their applied concentration.^{14b} Because most additives are applied to tobacco at very low concentrations (less than one-hundredth of one percent) the amounts that transfer to the smoke are minuscule. Other additives which are added to tobacco, such as the fruit juice extracts, also contain volatile flavor components that exhibit similar transfer and combustion behavior. Moreover, many of the juice extracts, i.e., prune juice and fig juice concentrate, used as cigarette flavoring agents comprise

2029063047

primarily sugars and simple carbohydrates (mainly glucose, fructose, xylose and sucrose) and, therefore, their contribution to mainstream smoke composition is expected to be similar to that of the sugars and carbohydrate components naturally found in tobacco.

SELECTION OF COMPONENTS FOR REVIEW

The selection of materials for inclusion in this report is based on a review of the literature on tobacco composition. A representative weight percent composition for a typical domestic blend (including additives) is summarized in Table 1.²⁹ Tobacco leaf is reported to contain over 2500 identified compounds.^{9,63}

The major additives used in tobacco products that contribute to mainstream smoke composition were selected for comparison to naturally-occurring tobacco components. Table 1 also summarizes the percent contribution of humectants, casing and flavor ingredients added to a typical domestic blend. Other major additives, such as urea and vanillin, were also included in the set of compounds selected for comparison. Table 2 summarizes the list of tobacco components and additives that are included in this review.

Table 3 summarizes cigarette mainstream smoke composition. Cigarette smoke is a dynamic aerosol having a discontinuous phase partially comprising particulate matter

2029063048

and a continuous phase composed in part of vapor and gas constituents. The whole smoke contains nearly 4,000 components.^{9, 32, 43}

APPROACHES FOR INVESTIGATING TOBACCO
INGREDIENTS AND COMPONENT PYROLYSIS PRODUCTS

Three basic methods are used to study the fate of materials subjected to cigarette combustion. These are: (1) addition of standard materials to the tobacco (enrichment) followed by determination of changes in mainstream smoke composition; (2) addition of radiolabeled materials to the tobacco followed by determination of their fates; and (3) study of materials using model pyrolysis test systems.

Each of the above methods can provide valuable information but each method suffers from certain drawbacks. For example, use of the first approach (addition of standard materials) can yield information which is generally appropriate for additives used at relatively high concentrations (e.g., sugars, humectants, etc.). However, the method is of limited value in quantifying the incremental contribution of an additive to individual smoke constituents where the additive is structurally identical or similar to that of a natural tobacco leaf component. In such cases, it is usually difficult to distinguish the additive's contribution to smoke composition from the natural background of smoke components. Also, particular care is needed when

2029063049

amounts of the test ingredient added for investigation are much higher than actual use levels. In such cases, cigarette combustion properties and therefore smoke composition will be altered and the results obtained can be invalid.

Additionally, if one adds large amounts of materials in an attempt to study "endogenous" tobacco constituents (e.g., sugars, amino acids, etc.), it is probable that the added materials will not be in the same matrix or state as the indigenous materials; again the experiments could be invalid.

Use of radiolabeled additives is an approach that can provide useful data. Radiolabeling experiments, however, are difficult and expensive to conduct and pyrolysis of radiolabeled materials added at higher than actual use levels may still yield misleading results. Furthermore, placement of the radiolabel is critical.

Studies using model test systems can give an indication of an additive's potential pyrolysis products under very specific experimental conditions. Such studies are relatively inexpensive to conduct and have the advantage of producing results more rapidly than other approaches. Because of the simplicity in experimental design, however, the studies do not accurately represent the complex thermal, atmospheric, and physical conditions and interactions of actual in-cigarette pyrolysis. For most non-volatile compounds there have to be significant discrepancies in the results obtained

2029063050

from laboratory pyrolysis and the types of compounds formed during cigarette combustion. For example, the rate of thermal energy transfer to the substrate in a burning cigarette and the rate at which primary pyrolysis products are removed from the hot zone are conditions that cannot be easily duplicated under laboratory pyrolysis conditions. Similarly, the presence of oxygen clearly plays a major role in the smoking process, yet pyrolysis in the laboratory is sure to be conducted under oxygenation conditions that differ significantly from those found in the smoking process. In addition, the presence of other substances in tobacco is known to modify the pyrolytic pathway, but model test systems do not reflect the interaction of substances that occur in actual in-cigarette pyrolysis.

EVALUATION OF NATURALLY OCCURRING TOBACCO LEAF COMPONENTS

Pyrolysis products of naturally occurring components of tobacco leaf are summarized below. The major structural classes of components are:

- Sugars
- Polysaccharides
- Phenolics
- Organic Acids and their salts
- Amino acids and proteins
- Waxes, lipids and oils

2029063051

- Nicotine

Sugars. Several sugars are natural components of tobacco leaf. Sugars can be found in flue-cured and oriental tobacco in amounts ranging up to 33 percent on a dry weight basis, but typically are present in amounts ranging between 5 to 25 percent. Air-cured tobacco such as burley and Maryland types have significantly lower total sugar levels in the maximum range of up to 0.2 percent on a dry weight basis in the cured leaf. The term "sugar" refers to a variety of different, but structurally-related chemical carbohydrate. Sugars that are natural components of tobacco include glucose, fructose, other hexoses, sucrose, other disaccharides, polysaccharides and pentoses.

Some of these same materials are also used as ingredients in the manufacture of cigarettes. Sugars are major constituents of tobacco casings and are used to reduce irritation and mellow the taste. Sugars also play an important role in achieving the necessary physical properties of cigarettes. They serve as plasticizers and humectants, thus contributing to the moisture-holding property, elasticity and flexibility of the cut tobacco strands.

Moreover, a number of tobacco additives and flavorings, including fruit juices, honey, molasses extract, corn and maple syrups, and caramel, largely comprise sugars and are expected to contribute to smoke composition in a

2029063052

manner similar to the sugars naturally found in the leaf. Because the contributions of tobacco leaf sugars to smoke components are well known, this information can be applied to evaluation of the pyrolysis products of potentially new additives composed of similar sugars.

Studies have shown that the major portion of the sugars in tobacco undergo in-cigarette combustion. Only glucose and sucrose are reported to transfer intact to mainstream smoke and even in these instances the transfer rate is very low (roughly 0.5% transfer rate).^{11, 12} A wide range of pyrolysis products are formed from the sugars. The major pyrolyzates are carbon dioxide, carbon monoxide, and water. Small amounts of furfural, 5-hydroxymethyl furfural, 5-methyl furfural, and furan have also been reported.^{14a, 18, 21, 31, 56, 65.} Other reported less concentrated products include other substituted furans, phenols (especially catechol), volatile aldehydes and ketones, levulinic acid and benzene and other acids.^{2, 11} It is worth noting that levulinic acid is commonly used as a flavorant in tobacco.

There is good agreement between cigarette enrichment studies, cigarette radiolabel studies and appropriately designed and conducted pyrolysis studies on sugar combustion processes. The studies show that the different types of sugars exhibit substantially similar pyrolysis behavior. Glucose, fructose, other hexoses and sucrose yield the same

2029063053

basic pyrolysis products but can differ in some of the relative amounts of individual pyrolyzates formed. There are some minor differences in pyrolysis behavior of the various sugars. Fructose produces more furfural than does glucose or sucrose.^{11, 55} Sucrose, reportedly, also produces some maltol.^{54a} Pentoses yield pyrolysis products similar to those from hexoses, but with one less carbon for substituted furans.²⁷

Polysaccharides. Several polysaccharides are natural components of tobacco leaf. These include cellulose, starch, pectins, dextrans, and pentosans. Several starches, pectins, and dextrans are also found in some of the complex carbohydrates that are added to tobacco. Certain of the polysaccharides, such as cellulose and the pentosans, are also major components of cigarette paper. None of these materials is reported to transfer intact to mainstream smoke. Maltodextrin, a simple polysaccharide comprising D-glucose units, is added to tobacco as a flavorant and also falls within this category. The main pyrolysis products from this group are water, carbon dioxide, carbon monoxide, phenols, organic acids, furan and some substituted furans.^{2, 14a, 42, 53, 55, 56, 60, 61} Other lesser pyrolysis products reported are benzene, toluene, volatile aldehydes and ketones and possibly trace amounts of polyaromatic hydrocarbons.^{51, 54a, 55} This product mix is similar to that from sugars but shows

2029063054

greater emphasis on phenols, aromatics and acids.³¹ Pyrolysis investigations of these materials indicate the following general trends:

- All materials studied in this group, except pentosans, yield the same basic pyrolysis products but differ in the relative amounts of individual components.
- Pentosans yield pyrolysis products similar to those from other materials in this group but with one less carbon for substituted furans, for example.
- Cigarette radiolabel and enrichment studies agree well with appropriate model system studies.

Phenolics. Several phenolics are natural components of tobacco leaf. These include chlorogenic acid (3-(3,4-dihydroxycinnamoyl)quinic acid), lignins, tannins and rutin. A number of the complex natural materials that are added to tobacco (e.g., nutmeg/mace, anise and anise oil, and star anise and star anise oil) comprise in part phenolic compounds. As a class the phenolics are characterized by the presence of a phenol component and a non-phenol component (often a sugar or sugar acid). None of these materials is reported to transfer intact to mainstream smoke. The main pyrolysis products reported for this class of materials are noted below:

- Chlorogenic acid yields catechol, catechol derivatives and quinic acid pyrolysis products (sugar breakdown products).^{2,18,44,52,69}

2029063055

- Lignins yield phenols, i.e., catechol plus some guaiacol (methyl catechol) and eugenol (4-allyl-2-methoxyphenol).^{18,49,55,56}
- Tannins yield catechols, other phenols, and sugar (usually glucose) breakdown products.^{18,44,59}
- Rutin yields products much like those from tannins.^{44,52,69}

Pyrolysis investigations of these materials indicate the following general trends:

- All members of the phenolic class pyrolyze to yield the phenol portion of the material.
- The non phenol portion of each material (often a sugar or sugar acid) yields pyrolysis products expected for their respective chemical classes.
- In-cigarette and appropriate model pyrolysis studies agree well with each other regarding contributions of these materials to mainstream smoke.

Organic Acids. Several organic acids, most notably lactic, oxalic, malic and citric acid are natural constituents of tobacco leaf. Collectively, organic acids may comprise as much as 9 percent of the leaf. A number of these organic acids also are used as additives to tobacco. Lactic acid is added to tobacco in small amounts (up to 0.4%) as a flavorant during the manufacture of cigarettes to give the tobacco a buttery, smooth taste. Citric acid is added to tobacco in the

2029063056

manufacture of cigarettes as a casing, at levels in the range of approximately one-hundredth of one percent. A number of other organic acid and organic acid salt additives, e.g., potassium sorbate, levulinic acid, potassium citrate and tartaric acid, are structurally analogous to the organic acids found in tobacco.

Radioisotope studies on malic, citric and oxalic acids show that only 15 percent of added organic acid transfers into the mainstream smoke, mostly as carbon monoxide, carbon dioxide and other acidic material.

The main pyrolysis products reported for each of these acids are:

- Some lactic acid transfers intact but the majority decomposes (primarily to carbon dioxide and ethanol). In addition some formic acid and acetaldehyde may be formed upon lactic acid pyrolysis.^{5,13}
- Oxalic acid may transfer intact through sublimation, but mainly it will yield carbon dioxide, carbon monoxide and formic acid upon pyrolysis.⁴⁰
- Malic acid may transfer intact, but mainly yields carbon dioxide, carbon monoxide, water, succinic acid, fumaric acid, ethanol and other lesser volatiles.^{13,40}
- Citric acid pyrolyzes in cigarettes to carbon dioxide, water and citraconic anhydride plus lesser volatiles.^{40,48}

2029063057

Pyrolysis investigations of these materials also indicate that:

- Organic acids will yield primarily carbon dioxide; those with alcoholic hydroxyl groups (i.e., citric, lactic, and malic acids) will also yield water.
- Each organic acid will also yield other predictable pyrolysis products besides carbon dioxide and water.
- Cigarette radiolabel studies agree well with appropriate model studies.

Amino Acids and Proteins. Many proteins and amino acids occur naturally in tobacco leaf. A number of amino acids (e.g., L-glutamine and L-asparagine monohydrate) and certain proteinaceous natural materials (e.g., carob bean) may be used as additives to tobacco. Amino acids are organic acids containing one or more basic amino groups (NH_2), one or more acidic carboxyl groups (COOH), and another group (R-group) that is the characteristic portion of the particular amino acid. Proteins are comprised of chains of amino acids connected by peptide linkages. A broad range of amino acids and representative proteins have been studied in this group. Pyrolysis investigations of these materials indicate that:

- Overall, a protein and a comparable mixture of amino acids will yield similar pyrolysis products.⁶²
- Particular amino acid R-groups give characteristic pyrolysis products.

2029063058

- Pyrolysis products from cigarette and model studies agree well.

The primary pyrolysis products of proteins and amino acids are reported to be ammonia, carbon dioxide, carbon monoxide and breakdown products of the remaining R-groups (for example phenolics will yield phenols, pyridines will yield pyridines, etc.)^{1,14a,18,47,57a,62} Other reported products include hydrogen cyanide and other nitriles plus unsaturated aldehydes and acids.^{15,25,26} Sulfur containing amino acids and proteins also produce some carbon disulfide.^{45a}

In addition to the above compounds, trace amounts of heterocyclic amines have been reported to form as a result of pyrolysis of proteins and amino acids. These include indole types (Trp-P-1, Trp-P-2), quinoxalines (MeIQx), quinolines (IQ), pyridines (PhIP), delta-azacarbolines (Glu-P-1, Glu-P-2) and amino-alpha-carbolines (A alpha C, MeA alpha C).^{30,35,36,37,68}

Waxes, Lipids and Oils. Many waxes, lipids and oils occur naturally in tobacco leaf. These include simple long chain esters, glyceryl esters, long chain fatty acids and long chain hydrocarbons. Methoprene, a well-studied insect growth regulator used in very small quantities prior to processing to protect stored tobacco from the cigarette beetle, is a member of this class of compounds. Pyrolysis investigations of these materials indicate that:

2029063059

- Esters and acids will generally yield carbon dioxide during pyrolysis.³⁸

- The hydrocarbon chains reportedly undergo scissions producing either lower molecular weight alkanes, alkenes and some aromatics if behind the fire cone with little or no oxygen or volatile aldehydes and ketones if appreciable oxygen is present.^{6,17,38}

- Some of these materials transfer intact into the mainstream smoke particulate phase.⁶⁷

Nicotine. Nicotine is reported to transfer intact into mainstream smoke at about 8-10% of blend concentration for a full flavor filtered cigarette.²⁰ Pyrolysis products of nicotine are reported to include mainly carbon dioxide and several pyridines, primarily pyridine, 3-methyl pyridine and 3-vinyl pyridine.^{20,58}

Appropriate nicotine pyrolysis model studies agree well with combustion results.^{28,58}

EVALUATION OF CIGARETTE ADDITIVES

Polyols. Polyols refer to polyhydric alcohols -- compounds containing two or more hydroxyl groups. Those having three hydroxyl groups (trihydric) are glycerols, those with more than three are called sugar alcohols. Several polyols are added to tobacco. These include propylene glycol, glycerol, and the hexahydric sugar alcohol, sorbitol.

2029063060

Maltitol, a polyhydric alcohol used as a humectant and a sweetener, also is a member of this class of materials.

- Propylene glycol has been used in cigarettes for thirty-five years as a humectant and as a vehicle for the addition of flavors. A typical cigarette may contain 1.5 to 15 mg of propylene glycol, or about 0.2 to 2 percent of the cigarette tobacco by weight. [recalculate based on new MULs] Pyrolysis studies have shown that only a small portion of the propylene glycol added to cigarettes (from 5 to 12.6 percent) is transferred intact to the mainstream smoke.³⁴ The small amount of pyrolysis that does occur yields mostly water, acetone, propanal and carbon dioxide.⁶⁴ Although minute quantities of benzo[a]pyrene have been detected in the pyrolysis of neat propylene glycol at 200°C in an airstream, the level is comparable to that obtained from pyrolyzing raw tobacco.

- Glycerol occurs naturally in tobacco. It also is used in the manufacture of cigarettes as a humectant (to aid in moisture retention) and as a processing aid, at levels up to 3.6 percent by weight. Pyrolysis studies have shown that 5% of the glycerol transfers intact to the smoke from filtered cigarettes. A major portion of the glycerol (roughly 30%) is retained in the butt and some 4 percent is retained in the tipping material.³⁴ Reported glycerol pyrolysis products include acetone, acetaldehyde, water and acrolein.^{7,64}

2029063061

- Sorbitol, a hexahydric sugar alcohol, yields the same general pyrolysis products as those obtained from pyrolysis of hexoses.^{22,23}

Other Single Materials. This group of additives includes menthol, triacetin, urea, propylparaben and polyvinyl acetate/alcohol. Pyrolysis studies on these materials provide the following information:

- Extensive cigarette testing reveals very little pyrolysis of menthol (usually <2%, primarily to menthane and menthone).^{24,41} Transfer to mainstream smoke ranges from 14-20%, depending on cigarette design. The distribution of ¹⁴C-menthol (uniformly labeled) and its combustion and pyrolytic products in cigarette smoke have been studied. The mainstream smoke contained 28.9% of the total activity, with 44.3% in the sidestream smoke, and 26.9% in the butt. The major ¹⁴C-menthol smoke product in the mainstream smoke was unchanged menthol (98.9%). Minor components that were identified in the mainstream smoke were carbon dioxide (0.1% of total mainstream activity), menthane (0.2%), and menthone (0.4% present as a contaminant of the labeled menthol). The additional 0.4% of the mainstream activity was located in unidentified peaks in the gas radiochromatograph.

Cigarettes were labeled with ¹⁴C-menthol, smoked, and the fate of the menthol was followed using an inverse isotope dilution assay. The mainstream smoke contained 41.87%

2029063062

of the radiolabel with 41.41% being in the solids and 0.46% in the gases (CO, CO₂, other gases). Unchanged menthol accounted for approximately 96% of the label in the mainstream solids. The sidestream smoke contained 37.56% of the label with 33.62% being in the solids and 3.94% in the gases (CO, CO₂, other gases). Unchanged menthol accounted for approximately 92% of the label in the sidestream solids. An additional 20.49% of the label was recovered in the butts and filters and 0.07% in the ashes. The author only investigated the amount of unchanged menthol present in the mainstream and sidestream solids and did not present data on any combustion products of menthol in these solids.

These reports may be contrasted with model test studies of the pyrolysis of pure menthol.^{57b} Menthol was pyrolyzed in horizontal quartz tubes packed with quartz chips under a stream of dry nitrogen at 600 and 860°C. Significant amounts of menthol were found to be pyrolyzed under these conditions -- 22% at 600° and 84% at 860°. Pyrolysis products included benzene, toluene, and styrene. In contrast, the studies described above which tracked the fate of ¹⁴C-labeled menthol added to tobacco found that well over 95% of the menthol remained in the smoke or the butt as unchanged, unpyrolyzed menthol. This suggests that pyrolytic studies of pure compounds are poor predictors of what actually occurs when the compound is burned in the cigarette. This is

2029063063

particularly true when test conditions are not comparable for temperature, heating rates, and ambient gas compositions within a burning cigarette.

- Triacetin transfers almost completely intact from the tobacco section. A small portion of triacetin on the tobacco section has been reported to pyrolyze to acetic acid, glycerol, acetone, acetin and diacetin.⁴⁶

- Urea is a natural component of tobacco, and is added to tobacco as a flavorant. A cigarette to which urea has been added may contain 2.2 to 5.4 mg of urea, or less than 1% of the cigarette tobacco by weight. Urea reportedly lowers cigarette gas phase delivery across the board but does not alter relative deliveries of HCN, nitrogen oxides, carbon monoxide or aldehydes.⁴⁶ Model studies report urea pyrolysis products to be water, ammonia and carbon dioxide.³³ Urea doesn't appear to transfer intact into cigarette smoke.

One study has reported that urea when added to tobacco reduces the amount of certain pyrolysis products in smoke: 43% decrease in benzo[a]pyrene; 15% decrease in CO; and 20% decrease in crude condensate.^{45b} In addition, the study indicated that urea impedes the formation of N-nitrosamines.

- Propylparaben (propyl-parahydroxy benzoate) is used in the manufacture of some reconstituted tobacco to prevent spoilage. Chromatographic (GC/MS) analysis of propylparaben

2029063064

pyrolysis indicates that at 315°C, the material distills without pyrolytic cleavage. At 590°C a very small amount of phenol is generated, but the vast majority of the compound transfers unchanged.

- Polyvinyl acetate is reported to yield acetic acid upon pyrolysis.^{19a} Polyvinyl acetate and polyvinyl alcohol are reported to yield common products including acetone, acetaldehyde, water and some crotonaldehyde.^{3,4,8}

Complex Natural Materials. Cocoa and licorice were reviewed in this group. The following results have been reported:

- Cocoa added to tobacco at levels of 1.3 percent reportedly results in a slight increase in tar, phenols, ammonia and selected other mainstream constituents.⁴⁶ Typically, cocoa used in cigarettes is about 12% butterfat, and as noted earlier, some of these materials can transfer intact into mainstream smoke particulate phase.⁶⁷ The composition of cocoa is well documented.^{16a} Cocoa powder contains more than 300 volatile compounds, many of which are produced during the roasting process. Many of those compounds also occur in smoke from cigarettes to which cocoa has not been added.^{16b}

Cocoa powder was pyrolyzed in a nitrogen atmosphere at temperatures from 370 to 750°C and the pyrolyzates were analyzed.^{54b} Cocoa powder produced palmitic and stearic acids

2029063065

on pyrolysis, particularly at 350 and 450°C, probably arising from thermal degradation of cocoa butter triglycerides. The author concluded that phenols derived from pyrolysis of cocoa powder should not significantly enhance the phenol content of tobacco smoke, although the fatty acid composition of smoke might be affected by triglyceride breakdown products.

The addition of cocoa to cigarettes, over a wide range of concentrations, has a minimal effect on the chemical composition of smoke. Tests on cigarettes cased with a variety of materials showed that, as the cocoa content of the casing mixture increased from 1.5 to 7.5 g/100 g of tobacco, the tar yield per cigarette increased from 19.2 mg to 19.7 mg; the benzo[a]pyrene from 10.3 nanograms to 11.6 nanograms; phenol from 45 micrograms to 48 micrograms; ammonia from 36 micrograms to 49.5 micrograms; and benzo[a]pyrene/TPM from 0.43 nanograms/mg to 0.48 nanograms/mg. Other components of smoke that were measured either decreased, remained the same, or were variable.

In another study, cocoa bean hulls were formed into thin sheets that were made into cigarettes. Levels of pyrene, fluoranthene, benzo[b]fluoranthene, and benzo[a]pyrene in smoke condensate from these cocoa "cigarettes" were not as high as the maximum levels of these compounds in tobacco cigarettes.^{19b}

2029063066

• Licorice is used in cigarettes both as a flavorant and as a casing material to smooth the harsh taste of certain kinds of tobacco. Most cigarette tobacco blends contain less than 1 percent licorice. Licorice contains from 6 to 14 percent of the tribasic acid glycyrrhizic acid, also known as glycyrrhizin. Besides glycyrrhizin, licorice contains sugars, starch, plant gums, resins, flavonoids, essential oils, inorganic salts and low levels of nitrogenous constituents such as proteins, amino acids, and nucleic acids.^{10,39} Licorice pyrolysis products reportedly include furans (from sugars), phenols (e.g., guaiacol) and glycerol plus a small amount of glycyrrhizic acid.^{50,66} The fate of glycyrrhizic acid and glycyrrhetinic acid during the smoking of cigarettes has been studied by adding these acids separately to cigarettes.⁵⁰ It was found that glycyrrhizic acid decomposed to glycyrrhetinic acid and was transferred in the mainstream smoke as such. When glycyrrhetinic acid itself was added to cigarettes, it was transferred intact to the mainstream smoke. In both cases, the amount of glycyrrhetinic acid found in the smoke condensate was small, and it was concluded that the glycyrrhizic acid in licorice root used for tobacco flavoring was mostly decomposed on smoking.

Several low level volatile materials occurring naturally in cocoa and licorice have been reported to transfer intact to smoke.^{16a,50,66}

2029063067

Processing Aids. Ethyl alcohol and carbon dioxide are used as processing aids in the manufacture of cigarettes. Because of their extremely high volatility and the manner in which they are used in the manufacturing process, virtually 100 percent of these materials is lost through volatilization and they are not incorporated into the final tobacco products.

Ethyl alcohol occurs naturally in tobacco smoke at a level of approximately 2 micrograms/cigarette. Although it also is used in cigarette manufacture as a solvent or carrier for flavor ingredients, essentially all of the added ethyl alcohol is volatilized during the manufacturing process, and the smoke contains virtually no added incremental level of ethyl alcohol.

Carbon dioxide is formed naturally during cigarette smoking, and accounts for roughly 13 percent of all mainstream smoke. The carbon dioxide used in the manufacture of cigarettes is lost through the volatilization processes and virtually none of it remains in the tobacco.

CONCLUSIONS

There are three primary approaches for investigating the contributions of tobacco leaf components and cigarette blend additives to cigarette mainstream smoke. Each has its unique strengths and limitations.

Despite these limitations, results from laboratory simulation studies generally agree with in-cigarette test

2029063068

results when test conditions are comparable for temperature, heating rates, and ambient gas compositions.

Materials sharing common chemical features yield similar pyrolysis products but relative amounts of individual components may differ. Thus, pyrolysis studies of cigarette additives and tobacco leaf components of similar chemical structure and/or composition yield similar pyrolysis products.

The approach of using compositional analogy for tobacco leaf components and additives provides a sound technical basis for predicting the potential pyrolysis products of proposed new tobacco additives.

Addition of volatile aromatic flavor substances is expected to impart specific flavor notes by virtue of their intact transfer into tobacco smoke. The concentrations of these ingredients, however, are extremely small and they generally are of similar chemical classes that occur naturally in tobacco and tobacco smoke.

While inclusion of non-volatile additives is expected to make small changes in the relative amounts of individual smoke components or classes of smoke components, these changes are not expected to significantly depart from the amounts or types of components generated from a range of additive free tobaccos or tobacco blends.

Attachment

2029063069

BIBLIOGRAPHY

1. Bryan, A. M. and Olafsson, P. G., Analysis of thermal decomposition patterns of aromatic and heterocyclic amino acids. *Anal. Lett.*, 2(10):505-513, 1969.
2. Carmella, S. G., Hecht, S. S., Tso, T. C., and Hoffmann, D., Roles of tobacco cellulose, sugars and chlorogenic acid as precursors to catechol in cigarette smoke. *J. Agric. Food Chem.*, 32(3):267-273, 1984.
3. Challinor, J. M., A pyrolysis-derivatization-gas chromatography technique for the structural elucidation of some synthetic polymers. *J. Anal. Appl. Pyrolysis*, 16(4):323-333, 1989.
4. Chytry, V., Obereigner, B. and Lim, D., Study of thermal decomposition of poly (vinyl chloride) type polymers with the use of model systems-III. Gas phase of acetates. *Eur. Polym. J.*, 9(6):493-500, 1973.
5. CRC Handbook of Food Additives. 2nd edition. Furia, T. E., (ed.), The Chemical Rubber Co., Cleveland, p. 237, 1972.
6. Davis, D. L., Waxes and lipids in leaf and their relationship to smoking quality and aroma. 30th Tobacco Chemists' Research Conference, Nashville, Tennessee: Recent Advances in Tobacco Science. 2:80-111, 1976.
7. DeSouza, J. E. and Scherbak, M., The effect of glycerol added to tobacco on the constituents of cigarette smoke. *Analyst*, 89:735-739, 1964.
8. Desreux, V. and Servotte, A., Thermal stability of polymers. Poly (vinyl acetate) and poly(vinyl alcohol). *Oberflaeche-Surface*, 11(10):268-278, 1970.
9. Dube, M. F. and Green, C. R., Methods of collection of smoke for analytical purposes. 36th Tobacco Chemists' Research Conference, Raleigh, North Carolina: Recent Advances in Tobacco Science. 8:42-102, 1982.
10. Frattini, C., Bicchi, C., Barettoni, C. and Nano, G. M., Volatile flavor components of licorice. *J. Agric. Food Chem.*, 25(6):1238-1241, 1977.

2029063070

11. Gager, F. L., Nedlock, J. W. and Martin, W. J., Tobacco additives and cigarette smoke. Part I. Transfer of D-glucose, sucrose and their degradation products to the smoke. Carbohyd. Res., 17:327-333, 1971.
12. Gager, F. L., Nedlock, J. W. and Martin, W. J., Tobacco additives and cigarette smoke. Part II. Organic, gas-phase products from D-glucose and sucrose. Carbohyd. Res., 17:335-339, 1971.
13. Geisinger, K. R., Jones, T. C. and Schmeltz, I., Further studies of pyrolytic products from some tobacco leaf acids. Tob. Sci., 14:89-90, 1970.
- 14a. Green, C. R., Some relationships between tobacco leaf and smoke composition, in Recent Advances in the Chemical Composition of Tobacco and Tobacco Smoke. The 173rd Amer. Chem. Soc. Meeting, New Orleans, Louisiana, 426-470, 1977.
- 14b. Green, J. D., Chalmers, J. and Kinnard, P. J., The Transfer of Tobacco Additives to Cigarette Smoke. Beitr. Tabaleforsch. Int., 14: 283-288, 1989.
15. Haidar, N. F., Patterson, J. M., Moors, M. and Smith, W. T. Jr., Effects of structure on pyrolysis gases from amino acids. J. Agric. Food Chem., 29(1):163-165, 1981.
- 16a. Harllee, G. and Leffingwell, J. C., Composition of casing materials: Cocoa, its constituents and their organoleptic properties. 32nd Tobacco Chemists' Research Conference, Montreal, Quebec: Paper 25, 1978.
- 16b. Harllee, G. and Leffingwell, J. C., Casing Materials -- Cocoa (Part I), Tobacco Int., 181(5):18-34, 40-52, 1979.
17. Higman, E. B., Schmeltz, I., Higman, H. C. and Chortyk, O. T., Studies on the thermal degradation of naturally occurring materials. Products from the pyrolysis of triglycerides at 400. J. Agric. Food Chem., 21(2):202-204, 1973.
18. Higman, E. B., Schmeltz, I. and Schlotzhauer, W. S., Products from the thermal degradation of some naturally occurring materials. J. Agric. Food Chem., 18:636-639, 1970.

2029063071

- 19a. Hiramatsu, K., Pyrolysis products of polymer materials studied by mass spectrometry. V. Mass spectrometry of pyrolysis products of polyamide (nylon 6) and poly(vinyl acetate). *Shitsuryo Bunseki*, 16(2):83-91, 1968.
- 19b. Hoffmann, D. et. al., Chemical Studies on Tobacco Smoke. XXXIX. On the Identification of Carcinogens, Tumor Promoters, and Co-carcinogens in Tobacco Smoke, in Wynder, E. L. et. al. (Ed.), Proceedings of the Third World Conference on Smoking and Health, June 2-5, 1975, Volume I, DHEW Publication No. (NIH) 76-1221, 1976, 125-145.
20. Houseman, T. H., The transfer of radiolabelled nicotine to cigarette smoke. 25th Tobacco Chemists' Research Conference, Louisville, Kentucky: Paper 9, 1971.
21. Houminer, Y. and Patai, S., Pyrolytic reactions of carbohydrates. Part II. Thermal decomposition of D-glucose. *Isr. J. Chem.*, 7(4):513-524, 1969.
22. Inagaki, H., Pyrolysis of sorbitol. Part 1. Identification of aldehydes. *Miyakonojo Kogyo Koto Semmon Gakko Kenkyu Hokoku*, 15:41-47, 1980.
23. Inagaki, H., Pyrolysis of sorbitol. Part 2. Identification of the lower boiling components of the pyrolyzate. *Miyakonojo Kogyo Koto Semmon Gakko Kenkyu Hokoku*, 17:25-34, 1982.
24. Jenkins, R. W. Jr., Newman, R. H. and Chavis, M. K., Cigarette smoke formation studies. II. Smoke distribution and mainstream pyrolytic composition of added 14 C-menthol(U). *Beitr. Tabakforsch.*, 5:299-301, 1970.
25. Johnson, W. R. and Kang, J. C., Mechanisms of hydrogen cyanide formation from the pyrolysis of amino acids and related compounds. *J. Org. Chem.*, 36(1):189-192, 1971.
26. Johnson, W. R., Nedlock, J. W. and Hale, R. W., Mechanisms of the pyrolysis of poly (amino acids). *Tob. Sci.*, 17:89-92, 1973.
27. Kaburaki, Y., Kobashi, U., Doihara, T. and Sugawara, S., Products of sugars pyrolysis. *Nippon Senbai Kosha Chuo Kenkyusho Kenkyu Hokoku*, 108:355-359, 1966.

2029063072

28. Kaburaki, Y., Sugawara, S., Kobashi, U. and Doihara, T., Composition of tobacco smoke. XIV. Formation of pyridines in pyrolysis of nicotine. Nippon Nogei Kagaku Kaishi, 44(5):224-231, 1970.
29. Kallianos, A. G., Phenolics and acids in leaf and their relationship to smoking quality and aroma. 30th Tobacco Chemists' Research Conference, Nashville, Tennessee: Recent Advances in Tobacco Science. 2:61-79, 1976.
30. Kanai, Y. Wada, O. and Manabe, S., Detection of carcinogenic glutamic acid pyrolysis products in cigarette smoke condensate. Carcinogenesis, 11(6):1001-1003, 1990.
31. Kato, K., Pyrolysis of cellulose. III. Comparative studies of the volatile compounds from pyrolyzates of cellulose and its related compounds. Agric. Biol. Chem., 31(6):657-663, 1967.
32. Keith, C. H. and Tesh, P. G., Measurement of total smoke issuing from a burning cigarette. Tob. Sci., 9:61-64, 1965.
33. Koryakin, A. G., Gal'perin, V. A., Sarbaev, A. N. and Finkel'shtein, A. I., Thermographic analysis of urea and products of its pyrolysis. Zh. Org. Khim., 7(5):972-977, 1971.
34. Laurene, A. H., Cundiff, R. H. and Greene, G. H., Determination of glycerol and propylene glycol in cigarette smoke. Tob. Sci., 9:1-4, 1965.
35. Manabe, S., Tohyama, K., Wada, O. and Aramaki, T., Detection of a carcinogen, 2-amino-1-methyl-6-phenylimidazo[4,5-b] pyridine (PhIP), in cigarette smoke condensate. Carcinogenesis, 12(10):1945-1947, 1991.
36. Manabe, S. and Wada, O., Carcinogenic tryptophan pyrolysis products in cigarette smoke condensate and cigarette smoke-polluted indoor air. Environ. Pollut., 64:121-132, 1990.
37. Manabe, S., Wada, O. and Kanai, Y., Simultaneous determination of amino-alpha-carbolines and amino-gamma-carbolines in cigarette smoke condensate by high-performance liquid chromatography. J. Chromatog., 529:125-133, 1990.

2029063073

38. Miyagawa, M., Pyrolysis of esters of saturated fatty acids. IV. Pyrolysis products of ethyl caproate. *Yukagaku*, 23(3):163-166, 1974.
39. Neiman, C., Licorice. *Adv. in Food Research*, 7:339-381, 1957.
40. Newell, M. P. and Best, F. W., The fate of oxalic, malic and citric acids in the smoke of cigarettes. 22nd Tobacco Chemists' Research Conference, Richmond, Virginia: Paper 25, 1968.
41. Newell, M. P., Latimer, P. H. and Haeefe, R. J., The fate of menthol in cigarette smoke. 22nd Tobacco Chemists' Research Conference, Richmond, Virginia: Paper 24, 1968.
42. Newell, M. P. and Best, F. W., Fate of compounds in the burning cigarette. Polymeric carbohydrates: pectins, starch, cellulose. 25th Tobacco Chemists' Research Conference, Louisville, Kentucky: Paper 11, 1971.
43. Norman, V., An overview of the vapor phase, semivolatile and nonvolatile components of cigarette smoke. 31st Tobacco Chemists' Research Conference, Greensboro, North Carolina: Recent Advances in Tobacco Science. 3:28-58, 1977.
44. Park, Jeon Woo, Study on the pyrolysis of polyphenols from tobacco by direct inlet/ms. *Yakhak Hoe Chi*, 26(2):123-128, 1982.
- 45a. Patterson, J. M., Shine, C. and Smith, W. T. Jr., Pyrolysis of some sulfur-containing amino acids at 850°C. *J. Agric. Food Chem.*, 24(5), 988-991, 1976.
- 45b. Pintaske, P., Additive for Smoking Tobacco, Ger. Pat. No. 2,942,544 (Apr. 30, 1981).
46. Private communications.
47. Ratcliff, M. A. Jr., Medley, E. E. and Simmonds, P. G., Pyrolysis of amino acids. Mechanistic considerations. *J. Org. Chem.*, 39(11):1481-1490, 1974.
48. Reiff, F. and Krueger, F., Citraconic anhydride, Ger. Pat. No. 1,283,229 (Nov. 21, 1968).

2029063074

49. Saiz-Jimenez, C. and DeLeeuw, J. W., Lignin pyrolysis products: Their structures and their significance as biomarkers. *Org. Geochem.*, 10(4-6):869-876, 1986.
50. Sakagami, H., Behavior of glycyrrhizic acid and glycyrrhetic acid added to tobacco on smoking -- Studies on the components of licorice root used for tobacco flavoring III. *Nippon Nogei Kagaku Kaishi*, 47(10):623-626, 1973; *Chem. Abstr.* 80, 93264d, 1974.
51. Sakuma, H., Kusama, M., Ishiguro, S., Shimojima, N. and Sugawara, S., Organic acid composition of cellulose cigarette smoke condensate. *Agric. Biol. Chem.*, 40(10):2021-2025, 1976.
52. Sakuma, H., Matsushima, S., Munakata, S. and Sugawara, S., Pyrolysis of chlorogenic acid and rutin. *Agric. Biol. Chem.*, 46(5):1311-1317, 1982.
53. Sakuma, H., Sachiko, M. and Sugawara, S., Volatile products of cellulose pyrolysis. *Agric. Biol. Chem.*, 45(2):443-451, 1981.
- 54a. Schlotzhauer, W. S., Arrendale, R. F. and Chortyk, O. T., The rapid pyrolytic characterization of tobacco leaf carbohydrate material. *Beitr. Tabakforsch. Int.*, 13:74-80, 1985.
- 54b. Schlotzhauer, W.S., Fatty Acids and Phenols from Pyrolysis of Cocoa Powder, a Tobacco Product Flavorant, *Tob. Sci.*, 22:1-2, 1978.
55. Schlotzhauer, W. S., Martin, R. M., Snook, M. E. and Williamson, R. E., Pyrolytic studies on the contribution of tobacco leaf constituents to the formation of smoke catechols. *J. Agric. Food Chem.*, 30(2):372-374, 1982.
56. Schlotzhauer, W. S., Schmeltz, I. and Hickey, L., Pyrolytic formation of phenols from some high molecular weight tobacco leaf constituents and related non-tobacco materials. *Tob. Sci.*, 11(8):31-34, 1967.
- 57a. Schmeltz, I., Schlotzhauer, W. S. and Higman, E. B., Characteristic products from pyrolysis of nitrogenous organic substances. *Beitr. Tabakforsch.*, 6:134-138, 1972.
- 57b. Schmeltz, I., Schlotzhauer, W.S., *Nature*, 219:370, 1968.

2029063075

58. Schmeltz, I., Wenger, A., Hoffmann, D. and Tso, T. C., Chemical studies on tobacco smoke. 63. On the fate of nicotine during pyrolysis and in a burning cigarette. *J. Agric. Food Chem.*, 27(3):602-608, 1979.
59. Schulman, G. P. and Simmonds, P. G., Thermal decomposition of aromatic and heterocyclic amino acids. *Chem. Commun.*, 17:1040-1042, 1968.
60. Shimada, M. and Deki, M., Thermal decomposition products of starch and its derivatives. *Kanzei Chuo Bunsekishoho*, 15:121-126, 1975.
61. Sjoberg, A. and Pyysalo, H., Identification of food thickeners by monitoring of their pyrolytic products. *J. Chromatogr.*, 319(1):90-98, 1985.
62. Smith, W. T. Jr., Harris, T. B. and Patterson, J. M., Pyrolysis of soybean protein and an amino acid mixture having the same amino acid composition. *J. Agric. Food Chem.*, 22(3):480-483, 1974.
63. Stedman, R. L., The chemical composition of tobacco and tobacco smoke. *Chem. Rev.*, 68:153-207, 1968.
64. Stein, Y. S. and Antal, M. J. Jr., A study of gas-phase pyrolysis of glycerol. *Jour. Anal. Appl. Pyrolysis*, 4(4):283-296, 1983.
65. Tomasik, P., Palasinski, M. and Wiejak, S., The thermal decomposition of mono-, di-, and oligo-saccharides. *Adv. Carbohydr. Chem. and Biochem.*, 47:272-278, 1989.
66. Vora, P. S. and Tuorto, R. M., The behavior of licorice components during the burning process of tobacco products. 38th Tobacco Chemists' Research Conference, Atlanta, Georgia: Paper 38, 1984.
67. Wakeham, H., The Chemistry of Tobacco and Tobacco Smoke. Schmeltz, I., (ed.), Plenum Press, New York, pp. 1-20, 1972.
68. Yamashita, M., Wakabayashi, K., Nagao, M., Sato, S., Yamaizumi, Z., Takahashi, M., Kinae, N., Tomita, I. and Sugimura, T., Detection of 2-amino-3-methyl-imidazo[4,5-f]quinoline in cigarette smoke condensate. *Jpn. J. Cancer Research. (Gann)*, 77:419-422, 1986.

2029063076

69. Zane, A. and Wender, S. H., Pyrolysis products of rutin, quercetin and chlorogenic acid. Tob. Sci., 7:21-23, 1963.

2029063077

TABLE 1

% Chemical Composition of a Representative U.S. Cigarette Blend
(Tobacco Plus Additive)

	<u>Total %</u> <u>By Weight</u>	<u>% From</u> <u>Tobacco</u>	<u>% From</u> <u>Additives</u>
Carbohydrates:			
Sugars	12.0	8.0	4.0
Cellulose	9.5	9.5	<0.1
Pectins	8.5	8.5	<0.1
Starch, pentosans	2.5	2.5	<0.1
Proteins	6.0	6.0	<0.1
Amino acids	2.5	2.5	<0.1
Organic acids	9.0	9.0	<0.1
Phenols	6.5	6.5	<0.1
Lignins	3.5	3.5	<0.1
Nicotine	2.0	2.0	----
Volatile bases	2.0	2.0	----
Waxes, resins	8.0	8.0	<0.1
Humectants & other additives	6.0	---	5.6*
Water	12.0	12.0	----
Inorganics	6.5	6.5	----
Others	3.5	3.5	----
Totals	100.0%	90.0%	10.0%

* Humectants & other additives

Humectants	3.5
Cocoa	0.9
Licorice	0.9
Menthol	0.2
Volatile flavorants	<0.1

Subtotal 5.6%

[Note: Ammonia (0.5) was removed from the list of humectants.
Does this affect the accuracy of this Table?]

2029063078

TABLE 2
Materials Included in this Review

<u>Sugars</u>	<u>Waxes, lipids and oils</u>
Glucose	Simple esters
Fructose	Glyceryl esters
Sucrose	Acids
Others	Hydrocarbons
<u>Polysaccharides</u>	<u>Polyols</u>
Cellulose	Propylene glycol
Starch	Glycerol
Pectins	Sorbitol
Dextrins	<u>Other single materials</u>
Pentosans	Menthol
<u>Phenolics</u>	Nicotine
Chlorogenic acid	Triacetin
Lignins	Urea
Tannins	Propylparaben
Others	Polyvinylacetate/alcohol
<u>Complex natural materials</u>	<u>Organic acids</u>
Cocoa	Lactic
Licorice	Maltic
<u>Amino acids and proteins</u>	Oxalic
[examples?]	Citric

2029063079

TABLE 3

Mainstream Smoke Composition for a Representative
U.S. Blend Full Taste Filtered Cigarette

<u>Whole Smoke</u>	<u>% By Weight</u>
N ₂	62
O ₂	13
CO ₂	12.5
CO	4
H ₂ O*	2
H ₂	1
Particulate phase**	4
Vapor phase***	<u>1.5</u>
	100.0

* From particulate + vapor phase

** No H₂O; 17.6 mg tar + 1.4 mg nicotine

*** No H₂O; less the above gases; ~6.8 mg.

Particulate and Vapor Phase Compositions

<u>Constituent</u> <u>or Class</u>	<u>mg In</u> <u>Particulate Phase</u>	<u>mg In</u> <u>Vapor Phase</u>	<u>mg Total</u>	<u>Number of</u> <u>Identified</u> <u>Compounds</u>
Hydrocarbons	3.15	3.04	6.19	760
Aldehydes	1.58	1.35	2.93	110
Ketones	0.90	0.81	1.71	520
Nitrogen heterocycles	0.79	0.14	0.93	920
Esters	0.79	0.07	0.86	480
Organic acids	2.93	0.10	3.03	230
Alcohols	1.80	0.14	1.94	380
Nicotine	1.35	---	1.35	1
Phenols	0.79	---	0.79	280
Nitriles	---	0.61	0.61	110
Others	4.84	0.49	5.33	Hundreds
Totals	18.9	6.8	25.7	~3800

2029063050